

CHAPTER 7 TECHNIQUES OF INTEGRATION

7.1 Integration by Parts (page 287)

Integration by parts is the reverse of the product rule. It changes $\int u \, dv$ into uv minus $\int v \, du$. In case $u = x$ and $dv = e^{2x} \, dx$, it changes $\int xe^{2x} \, dx$ to $\frac{1}{2}xe^{2x}$ minus $\int \frac{1}{2}e^{2x} \, dx$. The definite integral $\int_0^2 xe^{2x} \, dx$ becomes $\frac{3}{4}e^4$ minus $\frac{1}{4}$.

In choosing u and dv , the derivative of u and the integral of dv/dx should be as simple as possible. Normally $\ln x$ goes into u and e^x goes into v . Prime candidates are $u = x$ or x^2 and $v = \sin x$ or $\cos x$ or e^x . When $u = x^2$ we need two integrations by parts. For $\int \sin^{-1} x \, dx$, the choice $dv = dx$ leads to $x \sin^{-1} x$ minus $\int x \, dx / \sqrt{1 - x^2}$.

If U is the unit step function, $dU/dx = \delta$ is the unit delta function. The integral from $-A$ to A is $U(A) - U(-A) = 1$. The integral of $v(x)\delta(x)$ equals $v(0)$. The integral $\int_{-1}^1 \cos x \delta(x) \, dx$ equals 1. In engineering, the balance of forces $-dv/dx = f$ is multiplied by a displacement $u(x)$ and integrated to give a balance of work.

- $$\begin{array}{lll} 1 - x \cos x + \sin x + C & 3 - xe^{-x} - x + C & 5 x^2 \sin x + 2x \cos x - 2 \sin x + C \\ 7 \frac{1}{2}(2x+1) \ln(2x+1) + C & 9 \frac{1}{2}e^x(\sin x - \cos x) + C & 11 \frac{e^{ax}}{a^2+b^2}(a \sin bx - b \cos bx) + C \\ 13 \frac{x}{2}(\sin(\ln x) - \cos(\ln x)) + C & 15 x(\ln x)^2 - 2x \ln x + 2x + C & 17 x \sin^{-1} x + \sqrt{1-x^2} + C \\ 19 \frac{1}{2}(x^2+1) \tan^{-1} x - \frac{x}{2} + C & 21 x^3 \sin x + 3x^2 \cos x - 6x \sin x - 6 \cos x + C & \\ 23 e^x(x^3 - 3x^2 + 6x - 6) + C & 25 x \tan x + \ln(\cos x) + C & 27 -1 \quad 29 -\frac{3}{4}e^{-2} + \frac{1}{4} \quad 31 -2 \\ 33 3 \ln 10 - 6 + 2 \tan^{-1} 3 & 35 u = x^n, v = e^x & 37 u = x^n, v = \sin x \quad 39 u = (\ln x)^n, v = x \\ 41 u = x \sin x, v = e^x \rightarrow \int e^x \sin x \, dx \text{ in 9 and } -\int x \cos x \, e^x \, dx. \text{ Then } u = -x \cos x, v = e^x \rightarrow \int e^x \cos x \, dx \\ \text{in 10 and } -\int x \sin x \, e^x \, dx \text{ (move to left side): } \frac{e^x}{2}(x \sin x - x \cos x + \cos x). \text{ Also try } u = xe^x, v = -\cos x. \\ 43 \int \frac{1}{2}u \sin u \, du = \frac{1}{2}(\sin u - u \cos u) = \frac{1}{2}(\sin x^2 - x^2 \cos x^2); \text{ odd} & 45 3\text{-step function; } 3e^x \cdot \text{step function} \\ 49 0; x\delta(x)] - \int \delta(x) \, dx = -1; v(x)\delta(x)] - \int v(x)\delta(x) \, dx & 51 v(x) = \int_x^1 f(x) \, dx \\ 53 u(x) = \frac{1}{k} \int_0^x v(x) \, dx; \frac{1}{k}(\frac{x}{2} - \frac{x^3}{6}); \frac{x}{k} \text{ for } x \leq \frac{1}{2}, \frac{1}{k}(2x - x^2 - \frac{1}{4}) \text{ for } x \geq \frac{1}{2}; \frac{x}{k} \text{ for } x \leq \frac{1}{2}, \frac{1}{2k} \text{ for } x \geq \frac{1}{2}. \\ 55 u = x^2, v = -\cos x \rightarrow -x^2 \cos x - (2x) \sin x - \int 2 \sin x \, dx & 57 \text{ Compare 23} \\ 59 uw'|_0^1 - \int_0^1 u'w' - u'w|_0^1 + \int_0^1 u'w' = [uw' - u'w]|_0^1 \\ 61 \text{ No mistake: } e^x \cosh x - e^x \sinhx = 1 \text{ is part of the constant } C \end{array}$$

- $$\begin{array}{l} 2 uv - \int v \, du = x(\frac{1}{4}e^{4x}) - \int \frac{1}{4}e^{4x} \, dx = e^{4x}(\frac{x}{4} - \frac{1}{16}) + C \\ 4 uv - \int v \, du = x(\frac{1}{3} \sin 3x) - \int \frac{1}{3} \sin 3x \, dx = \frac{x}{3} \sin 3x + \frac{1}{9} \cos 3x + C \\ 6 uv - \int v \, du = (\ln x) \frac{x^2}{2} - \int \frac{x^2}{2} \frac{dx}{x} = \frac{x^2}{2} \ln x - \frac{x^2}{4} + C \\ 8 uv - \int v \, du = x^2(\frac{1}{4}e^{4x}) - \int (\frac{1}{4}e^{4x})2x \, dx = (\text{Problem 2}) e^{4x}(\frac{x^2}{4} - \frac{x}{8} + \frac{1}{32}) + C \\ 10 \int e^x \cos x \, dx = e^x \sin x - \int e^x \sin x \, dx. \text{ Another integration by parts produces } e^x(\sin x + \cos x) - \int e^x \cos x \, dx. \end{array}$$

Move the last integral to the left side and divide by 2: answer $\frac{1}{2}e^x(\sin x + \cos x) + C$.

- $$12 \text{ Not by parts. Substitute } u = x^2, du = 2x \, dx : \int \frac{1}{2}e^{-u} \, du = -\frac{1}{2}e^{-u} = -\frac{1}{2}e^{-x^2} + C.$$

- 14** $\int \cos(\ln x) dx = uv - \int v du = \cos(\ln x)x + \int x \sin(\ln x) \frac{1}{x} dx =$ again by parts gives $\cos(\ln x)x + \sin(\ln x)x - \int x \cos(\ln x) \frac{1}{x} dx$. Move the last integral to the left and divide by 2: answer $\frac{x}{2}(\cos(\ln x) + \sin(\ln x)) + C$.
- 16** $uv - \int v du = (\ln x) \frac{x^3}{3} - \int \frac{x^3}{3} \frac{dx}{x} = (\ln x) \frac{x^3}{3} - \frac{x^3}{9} + C$.
- 18** $uv - \int v du = \cos^{-1}(2x)x + \int x \frac{2 \frac{dx}{x}}{\sqrt{1-(2x)^2}} = x \cos^{-1}(2x) - \frac{1}{2}(1-4x^2)^{1/2} + C$.
- 20** $\int x^2 \sin x dx = x^2(-\cos x) + \int \cos x(2x dx) =$ again by parts gives $-x^2 \cos x + (\sin x)2x - \int \sin x(2 dx) =$ answer: $-x^2 \cos x + 2x \sin x + 2 \cos x + C$.
- 22** $uv - \int v du = x^3(-\cos x) + \int (\cos x)3x^2 dx =$ (use Problem 5) $= -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x + C$.
- 24** $uv - \int v du = \sec^{-1} x \left(\frac{x^2}{2} \right) - \int \frac{x^2}{2} \frac{dx}{|x|\sqrt{1-x^2}} = \frac{x^2}{2} \sec^{-1} x + \frac{1}{2}\sqrt{1-x^2} + C$.
- 26** $uv - \int v du = x \cosh x - \int \cosh x dx = x \cosh x - \sinh x + C$.
- 28** $\int_0^1 e^{\sqrt{x}} dx = \int_{u=0}^1 e^u (2u du) = 2e^u(u-1)|_0^1 = 2$. **30** $\ln(x^2) = 2 \ln x$; $\int_1^e 2 \ln x dx = [2(x \ln x - x)]_1^e = 2$.
- 32** $\int_{-\pi}^{\pi} x \sin x dx = [\sin x - x \cos x]_{-\pi}^{\pi} = 2\pi$.
- 34** $\int_0^{\pi/2} x^2 \sin x dx =$ (Problem 20) $[-x^2 \cos x + 2x \sin x + 2 \cos x]|_0^{\pi/2} = \pi - 2$.
- 36** $\int x^n e^{ax} dx = x^n \frac{e^{ax}}{a} - \frac{n}{a} \int x^{n-1} e^{ax} dx$. **38** $\int x^n \sin x dx = -x^n \cos x + n \int x^{n-1} \cos x dx$.
- 40** $\int x(\ln x)^n dx = (\ln x)^n \frac{x^2}{2} - \int \frac{x^2}{2} n(\ln x)^{n-1} \frac{dx}{x} = \frac{x^2}{2}(\ln x)^n - \frac{n}{2} \int x(\ln x)^{n-1} dx$.
- 42** Try $u = \tan^{-1} x$ and $dv = xe^x dx$ so $v = (x-1)e^x$. Then $\int v du = \int \frac{x-1}{1+x^2} e^x dx$. I believe this cannot be done in closed form; that is true for $\int \frac{e^x}{x} dx$.
- 44** (a) $e^0 = 1$; (b) $v(0)$ (c) 0 (limits do not enclose zero).
- 46** $\int_{-1}^1 \delta(2x) dx = \int_{u=-2}^2 \delta(u) \frac{du}{2} = \frac{1}{2}$. Apparently $\delta(2x)$ equals $\frac{1}{2}\delta(x)$; both are zero for $x \neq 0$.
- 48** $\int_0^1 \delta(x - \frac{1}{2}) dx = \int_{-1/2}^{1/2} \delta(u) du = 1$; $\int_0^1 e^x \delta(x - \frac{1}{2}) dx = \int_{-1/2}^{1/2} e^{u+\frac{1}{2}} \delta(u) du = e^{1/2}; \delta(x)\delta(x - \frac{1}{2}) = 0$.
- 50** $\int_{-1}^1 U(x) \frac{dU}{dx} dx =$ (directly) $[\frac{1}{2}(U(x))^2]|_0^1 = \frac{1}{2}$.
- 52** $-\frac{dv}{dx} = x$ gives $v = -\frac{x^2}{2} + C = -\frac{x^2}{2} + \frac{1}{2}$; $-\frac{dv}{dx} = U(x - \frac{1}{2})$ gives a change in slope at $x = \frac{1}{2}$:
 $v = C$ for $x \leq \frac{1}{2}$ and $v = C - (x - \frac{1}{2})$ for $x \geq \frac{1}{2}$; take $C = \frac{1}{2}$ to make $v(1) = 0$;
 $-\frac{dv}{dx} = \delta(x - \frac{1}{2})$ gives $v = C$ for $x < \frac{1}{2}$ and $v = C - 1$ for $x > \frac{1}{2}$; take $C = 1$ to make $v(1) = 0$.
- 54** $\frac{\Delta U}{\Delta x} = \frac{1}{\Delta x}$ over the interval from $x = -\Delta x$ to $x = 0$. Elsewhere $\Delta U = 0$. The area under the graph is $(\frac{1}{\Delta x})\Delta x = 1$. As $\Delta x \rightarrow 0$ the area is tall and thin. In the limit $\int \delta(x) dx = 1$.
- 56** $(-1)^n \int \frac{d^n u}{dx^n} v_{(n-1)} dx = (-1)^n \frac{d^n u}{dx^n} v_{(n)} + (-1)^{n+1} \int \frac{d^{n+1} u}{dx^{n+1}} v_{(n)} dx$.
- 58** $\int_0^x f'(t) dt = [uv]_0^x - \int_0^x v du = [f'(t)(t-x)]_0^x + \int_0^x (x-t) f''(t) dt = xf'(0) + \int_0^x (x-t) f''(t) dt$.
- 60** $A = \int_1^e \ln x dx = [x \ln x - x]_1^e = 1$ is the area under $y = \ln x$. $B = \int_0^1 e^y dy = e - 1$ is the area to the left of $y = \ln x$. Together the area of the rectangle is $1 + (e - 1) = e$.
- 62** The derivative is $C(ae^{ax} \cos bx - be^{ax} \sin bx) + D(ae^{ax} \sin bx + be^{ax} \cos bx)$. This equals $e^{ax} \cos bx$ if $Ca + Db = 1$ and $-Cb + Da = 0$. These two equations give $C = \frac{a}{a^2+b^2}$ and $D = \frac{b}{a^2+b^2}$. Knowing the correct form in advance seems easier than integrating.

7.2 Trigonometric Integrals (page 293)

To integrate $\sin^4 x \cos^3 x$, replace $\cos^2 x$ by $1 - \sin^2 x$. Then $(\sin^4 x - \sin^6 x) \cos x dx$ is $(u^4 - u^6) du$. In terms of $u = \sin x$ the integral is $\frac{1}{5}u^5 - \frac{1}{7}u^7$. This idea works for $\sin^m x \cos^n x$ if m or n is odd.

If both m and n are even, one method is integration by parts. For $\int \sin^4 x dx$, split off $dv = \sin x dx$.

Then $-\int v \, du$ is $\int 3 \sin^2 x \cos^2 x$. Replacing $\cos^2 x$ by $1 - \sin^2 x$ creates a new $\sin^4 x \, dx$ that combines with the original one. The result is a reduction to $\int \sin^2 x \, dx$, which is known to equal $\frac{1}{2}(x - \sin x \cos x)$.

The second method uses the double-angle formula $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$. Then $\sin^4 x$ involves $\cos^2 2x$. Another doubling comes from $\cos^2 2x = \frac{1}{2}(1 + \cos 4x)$. The integral contains the sine of $4x$.

To integrate $\sin 6x \cos 4x$, rewrite it as $\frac{1}{2} \sin 10x + \frac{1}{2} \sin 2x$. The integral is $-\frac{1}{20} \cos 10x - \frac{1}{4} \cos 2x$. The definite integral from 0 to 2π is zero. The product $\cos px \cos qx$ is written as $\frac{1}{2} \cos(p+q)x + \frac{1}{2} \cos(p-q)x$. Its integral is also zero, except if $p = q$ when the answer is π .

With $u = \tan x$, the integral of $\tan^n x \sec^2 x$ is $\frac{1}{10} \tan^{10} x$. Similarly $\int \sec^n x (\sec x \tan x \, dx) = \frac{1}{10} \sec^{10} x$. For the combination $\tan^n x \sec^n x$ we apply the identity $\tan^2 x = 1 + \sec^2 x$. After reduction we may need $\int \tan x \, dx = -\ln |\cos x|$ and $\int \sec x \, dx = \ln(\sec x + \tan x)$.

$$1 \int (1 - \cos^2 x) \sin x \, dx = -\cos x + \frac{1}{3} \cos^3 x + C \quad 3 \frac{1}{2} \sin^2 x + C$$

$$5 \int (1 - u^2)^2 u^2 (-du) = -\frac{1}{3} \cos^3 x + \frac{2}{5} \cos^5 x - \frac{1}{7} \cos^7 x + C \quad 7 \frac{2}{3} (\sin x)^{3/2} + C$$

$$9 \frac{1}{8} \int \sin^3 2x \, dx = \frac{1}{16} (-\cos 2x + \frac{1}{3} \cos^3 2x) + C \quad 11 \frac{\pi}{2} \quad 13 \frac{1}{3} (\frac{3x}{2} + \frac{\sin 6x}{4}) + C$$

$$15 x + C \quad 17 \frac{1}{6} \cos^5 x \sin x + \frac{5}{6} \int \cos^4 x \, dx; \text{ use equation (5)}$$

$$19 \int_0^{\pi/2} \cos^n x \, dx = \frac{n-1}{n} \int_0^{\pi/2} \cos^{n-2} x \, dx = \dots = \frac{n-1}{n} \frac{n-3}{n-2} \dots \frac{1}{2} \int_0^{\pi/2} dx$$

$$21 I = -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \cos^2 x \, dx = -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \, dx - (n-1)I.$$

$$\text{So } nI = -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \, dx.$$

$$23 0, +, 0, 0, 0, - \quad 25 -\frac{2}{3} \cos^3 x, 0 \quad 27 -\frac{1}{2} (\frac{\cos 2x}{2} + \frac{\cos 200x}{200}), 0 \quad 29 \frac{1}{2} (\frac{\sin 200x}{200} + \frac{\sin 2x}{2}), 0$$

$$31 -\frac{1}{2} \cos x, 0 \quad 33 \int_0^{\pi} x \sin x \, dx = \int_0^{\pi} A \sin^2 x \, dx \rightarrow A = 2 \quad 35 \text{ Sum} = \text{zero} = \frac{1}{2} \text{ (left + right)}$$

$$37 p \text{ is even} \quad 39 p - q \text{ is even} \quad 41 \sec x + C \quad 43 \frac{1}{3} \tan^3 x + C \quad 45 \frac{1}{3} \sec^3 x + C$$

$$47 \frac{1}{3} \tan^3 x - \tan x + x + C \quad 49 \ln |\sin x| + C \quad 51 \frac{1}{2 \cos^2 x} + C \quad 53 A = \sqrt{2}, -\sqrt{2} \sin(x + \frac{\pi}{4})$$

$$55 4\sqrt{2} \quad 57 \frac{1000}{\sqrt{3}} \quad 59 \frac{1 - \cos x + \sin x}{1 + \cos x + \sin x} + C \quad 61 p \text{ and } q \text{ are 10 and 1}$$

$$2 \int \cos^3 x \, dx = \int (1 - \sin^2 x) \cos x \, dx = \sin x - \frac{\sin^3 x}{3} + C$$

$$4 \int \cos^5 x \, dx = \int (1 - \sin^2 x)^2 \cos x \, dx = \int (1 - 2\sin^2 x + \sin^4 x) \cos x \, dx = \sin x - \frac{2}{3} \sin^3 x + \frac{1}{5} \sin^5 x + C$$

$$6 \int \sin^3 x \cos^3 x \, dx = \int \sin^3 x (1 - \sin^2 x) \cos x \, dx = \frac{1}{4} \sin^4 x - \frac{1}{6} \sin^6 x + C$$

$$8 \int \sqrt{\sin x} \cos^3 x \, dx = \int \sqrt{\sin x} (1 - \sin^2 x) \cos x \, dx = \frac{2}{3} (\sin x)^{3/2} - \frac{2}{7} (\sin x)^{7/2} + C$$

$$10 \int \sin^2 ax \cos ax \, dx = \frac{\sin^3 ax}{3a} + C \text{ and } \int \sin ax \cos ax \, dx = \frac{\sin^2 ax}{2a} + C$$

$$12 \int_0^{\pi} \sin^4 x \, dx = \int_0^{\pi} (\frac{1 - \cos 2x}{2})^2 \, dx = \frac{1}{4} \int_0^{\pi} (1 - 2\cos 2x + \frac{1 + \cos 4x}{2}) \, dx = [\frac{3x}{8} - \frac{\sin 2x}{4} + \frac{\sin 4x}{32}]_0^{\pi} = \frac{3\pi}{8}.$$

$$14 \int \sin^2 x \cos^2 x \, dx = \int \frac{1 - \cos 2x}{2} \frac{1 + \cos 2x}{2} \, dx = \int \frac{1 - \cos^2 2x}{4} \, dx = \int (\frac{1}{4} - \frac{1 + \cos 4x}{8}) \, dx = \frac{x}{8} - \frac{\sin 4x}{32} + C$$

$$16 \int \sin^2 x \cos^2 2x \, dx = \int \frac{1 - \cos 2x}{2} \cos^2 2x \, dx = \int (\frac{1 + \cos 4x}{4} - \frac{\cos 2x}{2} (1 - \sin^2 2x)) \, dx =$$

$$\frac{x}{4} + \frac{\sin 4x}{16} - \frac{\sin 2x}{4} + \frac{\sin^3 2x}{12} + C. \text{ This is a hard one.}$$

$$18 \text{ Equation (7) gives } \int_0^{\pi/2} \cos^n x \, dx = [\frac{\cos^{n-1} x \sin x}{n}]_0^{\pi/2} + \frac{n-1}{n} \int_0^{\pi/2} \cos^{n-2} x \, dx. \text{ The integrated term is zero}$$

because $\cos \frac{\pi}{2} = 0$ and $\sin 0 = 0$. The exception is $n = 1$, when the integral is $[\sin x]_0^{\pi/2} = 1$.

$$20 \text{ Problem 18 yields } \int_0^{\pi/2} \cos^n x \, dx = \frac{n-1}{n} \int_0^{\pi/2} \cos^{n-2} x \, dx = \frac{n-1}{n} \frac{n-3}{n-2} \int_0^{\pi/2} \cos^{n-4} x \, dx. \text{ For odd } n \text{ this}$$

continues to $\frac{n-1}{n} \frac{n-3}{n-2} \dots \frac{2}{3}$, times $\int_0^{\pi/2} \cos x dx = 1$. Writing from low to high this is $\frac{2}{3} \frac{4}{5} \dots \frac{n-1}{n}$.

22 $\int_0^\pi \cos x dx = 0$ because the positive area from 0 to $\frac{\pi}{2}$ is balanced by the negative area from $\frac{\pi}{2}$ to π . This is true for any odd power $n = 1, 3, 5, \dots$ (For even powers $\cos^n x$ is always positive). The substitution $u = \pi - x$ and $du = -dx$ gives $\int_0^\pi \cos^n x dx = -\int_\pi^0 \cos^n(\pi - u) du = \int_0^\pi (-1)^n \cos^n u du$. So if n is odd, the integral equals minus the integral and must be zero.

24 $(\sin x)(\sin x) = -\frac{1}{2} \cos(1+1)x + \frac{1}{2} \cos(1-1)x$ is the double angle formula $\sin^2 x = \frac{1-\cos 2x}{2}$; $(\cos 2x)(\cos x) = \frac{1}{2} \cos(2+1)x + \frac{1}{2} \cos(2-1)x = \frac{\cos 3x + \cos x}{2}$. To derive equation (9), subtract $\cos(s+t) = \cos s \cos t - \sin s \sin t$ from $\cos(s-t) = \cos s \cos t + \sin s \sin t$. Divide by 2. Then set $s = px$ and $t = qx$.

$$26 \int_0^\pi \sin 3x \sin 5x dx = \int_0^\pi \frac{-\cos 8x + \cos 2x}{2} dx = \left[\frac{-\sin 8x}{16} + \frac{\sin 2x}{4} \right]_0^\pi = 0.$$

$$28 \int_{-\pi}^\pi \cos^2 3x dx = \int_{-\pi}^\pi \frac{1+\cos 6x}{2} dx = \left[\frac{x}{2} + \frac{\sin 6x}{12} \right]_{-\pi}^\pi = \pi.$$

$$30 \int_0^{2\pi} \sin x \sin 2x \sin 3x dx = \int_0^{2\pi} \sin 2x \left(\frac{-\cos 4x + \cos 2x}{2} \right) dx = \int_0^{2\pi} \sin 2x \left(\frac{1-2\cos^2 2x + \cos 2x}{2} \right) dx = \left[-\frac{\cos 2x}{4} + \frac{\cos^3 2x}{6} - \frac{\cos^2 2x}{8} \right]_0^{2\pi} = 0. \text{ Note: The integral has other forms.}$$

$$32 \int_0^\pi x \cos x dx = [x \sin x]_0^\pi - \int_0^\pi \sin x dx = [x \sin x + \cos x]_0^\pi = -2.$$

$$34 \int_0^\pi 1 \sin 3x dx = \int_0^\pi (A \sin x + B \sin 2x + C \sin 3x + \dots) \sin 3x dx \text{ reduces to } [-\frac{\cos 3x}{3}]_0^\pi = 0 + 0 + C \int_0^\pi \sin^2 3x dx. \text{ Then } \frac{2}{3} = C \left(\frac{\pi}{2} \right) \text{ and } C = \frac{4}{3\pi}.$$

36 The square wave is -1 and 1 periodically. To find A , multiply the series by $\sin x$ and integrate from 0 to π : $\int_0^\pi 1 \sin x dx = \int_0^\pi (A \sin x + \dots) \sin x dx$ yields $2 = A \left(\frac{\pi}{2} \right)$ and $A = \frac{4}{\pi}$. To find B , multiply the series by $\sin 2x$ and integrate: $\int_0^\pi 1 \sin 2x dx = \int_0^\pi (A \sin x + B \sin 2x + \dots) \sin 2x dx$ yields $0 = B \int_0^\pi \sin^2 2x dx$ and $B = 0$.

$$38 \int_0^\pi \cos qx dx = [\frac{\sin qx}{q}]_0^\pi = \frac{\sin q\pi}{q} \text{ which is zero if } q \text{ is any nonzero integer.}$$

$$40 \text{ "Always zero" means for positive integers } p \neq q. \text{ Then } \int_0^\pi \sin px \sin qx dx = \int_0^\pi \frac{-\cos(p+q)x + \cos(p-q)x}{2} dx = \left[\frac{-\sin(p+q)x}{2(p+q)} + \frac{\sin(p-q)x}{2(p-q)} \right]_0^\pi = 0.$$

$$42 \int \tan 5x dx = \int \frac{\sin 5x}{\cos 5x} dx = -\frac{1}{5} \ln |\cos 5x| \text{ (set } u = \cos 5x \text{ to find } \int \frac{du}{5u}).$$

44 First by substituting for $\tan^2 x$: $\int \tan^2 x \sec x dx = \int \sec^3 x dx - \int \sec x dx$. Use Problem 62 to integrate $\sec^3 x$: final answer $\frac{1}{2} (\sec x \tan x - \ln |\sec x + \tan x|) + C$. Second method from line 1 of Example 11: $\int \tan^2 x \sec x dx = \sec x \tan x - \int \sec^3 x dx$. Same final answer.

$$46 \int \sec^4 x dx = \int \sec^2 x (1 + \tan^2 x) dx = \tan x + \frac{\tan^3 x}{3} + C$$

$$48 \int \tan^5 x dx = \int (\sec^2 x - 1) \tan^3 x dx = \frac{\tan^4 x}{4} - \int \tan^3 x dx = \frac{\tan^4 x}{4} - \int (\sec^2 x - 1) \tan x dx = \frac{\tan^4 x}{4} - \frac{\tan^2 x}{2} - \ln |\cos x| + C$$

50 OK to write down $\ln |\csc x - \cot x|$ or $-\ln |\csc x + \cot x|$. For variety set $u = \frac{\pi}{2} - x$ and integrate $-\int \sec u du$.

52 This should have an asterisk! $\int \frac{\sin^6 x}{\cos^5 x} dx = \int \frac{(1-\cos^2 x)^3}{\cos^5 x} dx = \int (\sec^3 x - 3 \sec x + 3 \cos x - \cos^3 x) dx =$ use Example 11 = Problem 62 for $\int \sec^3 x dx$ and change $\int \cos^3 x dx$ to $\int (1 - \sin^2 x) \cos x dx$.

Final answer $\frac{\sec x \tan x}{2} - \frac{5}{2} \ln |\sec x + \tan x| + 2 \sin x + \frac{\sin^3 x}{3} + C$.

$$54 A = 2 : 2 \cos(x + \frac{\pi}{3}) = 2 \cos x \cos \frac{\pi}{3} - 2 \sin x \sin \frac{\pi}{3} = \cos x - \sqrt{3} \sin x. \text{ Therefore } \int \frac{dx}{(\cos x - \sqrt{3} \sin x)^2} = \int \frac{dx}{4 \cos^2(x + \frac{\pi}{3})} = \frac{1}{4} \tan(x + \frac{\pi}{3}) + C.$$

56 Expand $\cos(x - \alpha) = \cos x \cos \alpha + \sin x \sin \alpha$, multiply by $\sqrt{a^2 + b^2}$, and match with $a \cos x + b \sin x$.

Then $\cos \alpha = \frac{a}{\sqrt{a^2 + b^2}}$ and $\sin \alpha = \frac{b}{\sqrt{a^2 + b^2}}$ is correct if $\tan \alpha = \frac{b}{a}$ (the right triangle has sides a and b).

58 When lengths are scaled by $\sec x$, area is scaled by $\sec^2 x$. The area from the equator to latitude x is then proportional to $\int \sec^2 x dx = \tan x$.

60 The graphs of $\sin^2 x$ and $\cos^2 x$ obviously give equal areas between 0 and $\frac{\pi}{2}$ and between $\frac{\pi}{2}$ and π . The areas add to $\int_0^\pi 1 dx = \pi$ so each area is $\frac{\pi}{2}$.

- 62 Example 11 ends with $2 \int \sec^3 x \, dx = \sec x \tan x + \int \sec x \, dx$. Divide by 2 to find $\int \sec^3 x \, dx = \frac{1}{2}(\sec x \tan x + \ln|\sec x + \tan x|) + C$.

7.3 Trigonometric Substitutions (page 299)

The function $\sqrt{1 - x^2}$ suggests the substitution $x = \sin \theta$. The square root becomes $\cos \theta$ and dx changes to $\cos \theta \, d\theta$. The integral $\int (1 - x^2)^{3/2} \, dx$ becomes $\int \cos^4 \theta \, d\theta$. The interval $\frac{1}{2} \leq x \leq 1$ changes to $\frac{\pi}{6} \leq \theta \leq \frac{\pi}{2}$.

For $\sqrt{a^2 - x^2}$ the substitution is $x = a \sin \theta$ with $dx = a \cos \theta \, d\theta$. For $x^2 - a^2$ we use $x = a \sec \theta$ with $dx = a \sec \theta \tan \theta \, d\theta$. (Insert: For $x^2 + a^2$ use $x = a \tan \theta$). Then $\int dx/(1 + x^2)$ becomes $\int d\theta$, because $1 + \tan^2 \theta = \sec^2 \theta$. The answer is $\theta = \tan^{-1} x$. We already knew that $\frac{1}{1+x^2}$ is the derivative of $\tan^{-1} x$.

The quadratic $x^2 + 2bx + c$ contains a linear term $2bx$. To remove it we complete the square. This gives $(x+b)^2 + C$ with $C = c - b^2$. The example $x^2 + 4x + 9$ becomes $(x+2)^2 + 5$. Then $u = x+2$. In case x^2 enters with a minus sign, $-x^2 + 4x + 9$ becomes $-(x-2)^2 + 13$. When the quadratic contains $4x^2$, start by factoring out 4.

- 1 $x = 2 \sin \theta; \int d\theta = \sin^{-1} \frac{x}{2} + C$
- 3 $x = 2 \sin \theta; \int 4 \cos^2 \theta \, d\theta = 2 \sin^{-1} \frac{x}{2} + x \sqrt{1 - \frac{x^2}{4}} + C$
- 5 $x = \sin \theta; \int \sin^2 \theta \, d\theta = \frac{1}{2} \sin^{-1} x - \frac{1}{2} x \sqrt{1 - x^2} + C$
- 7 $x = \tan \theta; \int \cos^2 \theta \, d\theta = \frac{1}{2} \tan^{-1} x + \frac{1}{2} \frac{x}{1+x^2} + C$
- 9 $x = 5 \sec \theta; \int 5(\sec^2 \theta - 1) \, d\theta = \sqrt{x^2 - 25} - 5 \sec^{-1} \frac{x}{5} + C$
- 11 $x = \sec \theta; \int \cos \theta \, d\theta = \frac{\sqrt{x^2 - 1}}{x} + C$
- 13 $x = \tan \theta; \int \cos \theta \, d\theta = \frac{x}{\sqrt{1+x^2}} + C$
- 15 $x = 3 \sec \theta; \int \frac{\cos \theta \, d\theta}{9 \sin^2 \theta} = \frac{-1}{9 \sin \theta} + C = \frac{-x}{9\sqrt{x^2 - 9}} + C$
- 17 $x = \sec \theta; \int \sec^3 \theta \, d\theta = \frac{1}{2} \sec \theta \tan \theta + \frac{1}{2} \ln(\sec \theta + \tan \theta) + C = \frac{1}{2} x \sqrt{x^2 - 1} + \frac{1}{2} \ln(x + \sqrt{x^2 - 1}) + C$
- 19 $x = \tan \theta; \int \frac{\cos \theta \, d\theta}{\sin^2 \theta} = -\frac{1}{\sin \theta} + C = \frac{-\sqrt{x^2 + 1}}{x} + C$
- 21 $\int \frac{-\sin \theta \, d\theta}{\sin \theta} = -\theta + C = -\cos^{-1} x + C$; with $C = \frac{\pi}{2}$ this is $\sin^{-1} x$
- 23 $\int \frac{\tan \theta \sec^2 \theta \, d\theta}{\sec^2 \theta} = -\ln(\cos \theta) + C = \ln \sqrt{x^2 + 1} + C$ which is $\frac{1}{2} \ln(x^2 + 1) + C$
- 25 $x = a \sin \theta; \int_{-\pi/2}^{\pi/2} a^2 \cos^2 \theta \, d\theta = \frac{a^2 \pi}{2}$ = area of semicircle
- 27 $\sin^{-1} x |_{-5}^1 = \frac{\pi}{2} - \frac{\pi}{6} = \frac{\pi}{3}$
- 29 Like Example 6: $x = \sin \theta$ with $\theta = \frac{\pi}{2}$ when $x = \infty$, $\theta = \frac{\pi}{3}$ when $x = 2$, $\int_{\pi/3}^{\pi/2} \frac{\cos \theta \, d\theta}{\sin^2 \theta} = -1 + \frac{2}{\sqrt{3}}$
- 31 $x = 3 \tan \theta; \int_{-\pi/2}^{\pi/2} \frac{3 \sec^2 \theta \, d\theta}{9 \sec^2 \theta} = \frac{\theta}{3} |_{-\pi/2}^{\pi/2} = \frac{\pi}{3}$
- 33 $\int \frac{x^{n+1} + x^{n-1}}{x^2 + 1} \, dx = \int x^{n-1} \, dx = \frac{x^n}{n}$
- 35 $x = \sec \theta; \frac{1}{2}(e^f + e^{-f}) = \frac{1}{2}(x + \sqrt{x^2 - 1} + \frac{1}{x + \sqrt{x^2 - 1}}) = \frac{1}{2}(x + \sqrt{x^2 - 1} + x - \sqrt{x^2 - 1}) = x$
- 37 $x = \cosh \theta; \int d\theta = \cosh^{-1} x + C$
- 39 $x = \cosh \theta; \int \sinh^2 \theta \, d\theta = \frac{1}{2}(\sinh \theta \cosh \theta - \theta) + C = \frac{1}{2}x \sqrt{x^2 - 1} - \frac{1}{2} \ln(x + \sqrt{x^2 - 1}) + C$
- 41 $x = \tanh \theta; \int d\theta = \tanh^{-1} x + C$
- 43 $(x-2)^2 + 4$
- 45 $(x-3)^2 - 9$
- 47 $(x+1)^2$
- 49 $u = x-2, \int \frac{du}{u^2+4} = \frac{1}{2} \tan^{-1} \frac{u}{2} = \frac{1}{2} \tan^{-1} \left(\frac{x-2}{2} \right) + C; u = x-3, \int \frac{du}{u^2-9} = \frac{1}{6} \ln \frac{u-3}{u+3} = \frac{1}{6} \ln \frac{x-6}{x} + C;$
 $u = x+1, \int \frac{du}{u^2} = \frac{-1}{u} = \frac{-1}{x+1} + C$
- 51 $u = x+b; \int \frac{du}{u^2-b^2+c}$ uses $u = a \sec \theta$ if $b^2 > c$, $u = a \tan \theta$ if $b^2 < c$, equals $-\frac{1}{u} = \frac{-1}{x+b}$ if $b^2 = c$

53 $\cos \theta$ is negative ($-\sqrt{1-x^2}$) from $\frac{\pi}{2}$ to $\frac{3\pi}{2}$; then $\int_0^1 -\int_1^{-1} + \int_{-1}^0 \sqrt{1-x^2} dx = \pi =$ area of unit circle

55 Divide y by 4, multiply dx by 4, same $\int y dx$

57 No $\sin^{-1} x$ for $x > 1$; the square root is imaginary. All correct with complex numbers.

$$\mathbf{2} \quad x = a \sec \theta, x^2 - a^2 = a^2 \tan^2 \theta, \int \frac{dx}{\sqrt{x^2 - a^2}} = \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \ln |\sec \theta + \tan \theta| = \ln \left| \frac{x}{a} + \sqrt{\frac{x^2}{a^2} - 1} \right| + C$$

$$\mathbf{4} \quad x = \frac{1}{3} \tan \theta, 1 + 9x^2 = \sec^2 \theta, \int \frac{dx}{1+9x^2} = \int \frac{\frac{1}{3} \sec^2 \theta d\theta}{\sec^2 \theta} = \frac{\theta}{3} = \frac{1}{3} \tan^{-1} 3x + C.$$

$$\mathbf{6} \quad x = \sin \theta, \int \frac{dx}{x^2 \sqrt{1-x^2}} = \int \frac{\cos \theta d\theta}{\sin^2 \theta \cos \theta} = -\cot \theta = -\frac{\sqrt{1-x^2}}{x} + C$$

8 $x = a \tan \theta, x^2 + a^2 = a^2 \sec^2 \theta, \int \sqrt{x^2 + a^2} dx = \int a^2 \sec^3 \theta d\theta =$ use Problem 62 above:

$$\frac{a^2}{2} (\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|) = \frac{1}{2} x \sqrt{x^2 + a^2} + \frac{a^2}{2} \ln \left| \frac{x}{a} + \sqrt{\frac{x^2 + a^2}{a^2}} \right| + C$$

$$\mathbf{10} \quad x = 3 \sin \theta, 9 - x^2 = 9 \cos^2 \theta, \int \frac{x^3 dx}{\sqrt{9-x^2}} = \int \frac{27 \sin^3 \theta (3 \cos \theta d\theta)}{3 \cos \theta} = \int 27(1 - \cos^2 \theta) \sin \theta d\theta = -27 \cos \theta + 9 \cos^3 \theta = -27(1 - \frac{x^2}{9})^{1/2} + 9(1 - \frac{x^2}{9})^{3/2} + C$$

$$\mathbf{12} \quad \text{Write } \sqrt{x^6 - x^8} = x^3 \sqrt{1 - x^2} \text{ and set } x = \sin \theta : \int \sqrt{x^6 - x^8} dx = \int \sin^3 \theta \cos \theta (\cos \theta d\theta) = \int \sin \theta (\cos^2 \theta - \cos^4 \theta) d\theta = -\frac{\cos^3 \theta}{3} + \frac{\cos^5 \theta}{5} = -\frac{1}{3}(1 - x^2)^{3/2} + \frac{1}{5}(1 - x^2)^{5/2} + C$$

$$\mathbf{14} \quad x = \sin \theta, \int \frac{dx}{(1-x^2)^{3/2}} = \int \frac{\cos \theta d\theta}{\cos^3 \theta} = \tan \theta + C = \frac{x}{\sqrt{1-x^2}} + C.$$

$$\mathbf{16} \quad x = \tan \theta, \int \frac{\sqrt{1+x^2} dx}{x} = \int \frac{\sec \theta \sec^2 \theta d\theta}{\tan \theta} = \int \frac{\sec \theta (1+\tan^2 \theta) d\theta}{\tan \theta} = \int (\csc \theta + \sec \theta \tan \theta) d\theta = \ln |\csc \theta - \cot \theta| + \sec \theta = \ln \left| \frac{\sqrt{1+x^2}}{x} - \frac{1}{x} \right| + \sqrt{1+x^2} + C.$$

$$\mathbf{18} \quad x = 2 \tan \theta, x^2 + 4 = 4 \sec^2 \theta, \int \frac{x^2 dx}{x^2 + 4} = \int \frac{4 \tan^2 \theta}{4 \sec^2 \theta} 2 \sec^2 \theta d\theta = \int 2 \tan^2 \theta d\theta = \int 2(\sec^2 \theta - 1) d\theta = 2 \tan \theta - 2\theta = x - 2 \tan^{-1} \frac{x}{2} + C.$$

$$\mathbf{20} \quad x = \tan \theta, 1 + x^2 = \sec^2 \theta, \int \frac{x^2 dx}{\sqrt{1+x^2}} = \int \frac{\tan^2 \theta}{\sec \theta} \sec^2 \theta d\theta = \int \tan^2 \theta \sec \theta d\theta = (\text{use Problem 44 above}) \frac{1}{2} (\sec \theta \tan \theta - \ln |\sec \theta + \tan \theta|) = \frac{1}{2} (x \sqrt{1+x^2} - \ln |\sqrt{1+x^2} + x|) + C.$$

$$\mathbf{22} \quad x = \sec \theta : \int \frac{dx}{x \sqrt{x^2 - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{\sec \theta \tan \theta} = \theta + C = \sec^{-1} x + C. \text{ For } x = \csc \theta \text{ the integral is } \int \frac{-\csc \theta \cot \theta d\theta}{\csc \theta \cot \theta} = -\theta + C = -\csc^{-1} x + C^*. \text{ Both answers are right: } \sec^{-1} x + \csc^{-1} x = \text{sum of complementary angles in Section 4.4} = \frac{\pi}{2} \text{ so the arbitrary constant } C^* \text{ is } C - \frac{\pi}{2}.$$

$$\mathbf{24} \quad \text{Set } x^2 = \sec \theta \text{ and } x^4 - 1 = \tan^2 \theta \text{ and } 2x dx = \sec \theta \tan \theta d\theta. \text{ Then } \int \frac{2x dx}{2x^2 \sqrt{x^4 - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{2 \sec \theta \tan \theta} = \frac{\theta}{2} = \frac{1}{2} \sec^{-1}(x^2).$$

$$\mathbf{26} \quad x = \sin \theta : \int_{-1}^1 (1 - x^2)^{3/2} dx = \int_{-\pi/2}^{\pi/2} \cos^3 \theta (\cos \theta d\theta) = 2 \int_0^{\pi/2} \cos^4 \theta d\theta = (\text{Problem 19 of Section 7.2}) 2\left(\frac{1}{2}\right)\left(\frac{3}{4}\right)\left(\frac{\pi}{2}\right) = \frac{3\pi}{8}.$$

$$\mathbf{28} \quad x = \sec \theta : \int_1^4 \frac{dx}{\sqrt{x^2 - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{\tan \theta} = \ln |\sec \theta + \tan \theta| = [\ln |x + \sqrt{x^2 - 1}|]_1^4 = \ln (4 + \sqrt{15}).$$

$$\mathbf{30} \quad \int_{-1}^1 \frac{x dx}{x^2 + 1} = [\frac{1}{2} \ln(x^2 + 1)]_{-1}^1 = 0 \text{ (odd function integrated from } -1 \text{ to } 1).$$

32 First use geometry: $\int_{1/2}^1 \sqrt{1-x^2} dx =$ half the area of the unit circle beyond $x = \frac{1}{2}$ which breaks into

$$\frac{1}{2} (120^\circ \text{ wedge minus } 120^\circ \text{ triangle}) = \frac{1}{2} \left(\frac{\pi}{3} - \frac{1}{2} \cdot \frac{1}{2} \cdot 2 \sqrt{1 - (\frac{1}{2})^2} \right) = \frac{\pi}{6} - \frac{\sqrt{3}}{8}.$$

Check by integration: $\int_{1/2}^1 \sqrt{1-x^2} dx = [\frac{1}{2} (x \sqrt{1-x^2} + \sin^{-1} x)]_{1/2}^1 = \frac{1}{2} \left(\frac{\pi}{2} - \frac{1}{2} \frac{\sqrt{3}}{2} - \frac{\pi}{6} \right) = \frac{\pi}{6} - \frac{\sqrt{3}}{8}.$

$$\mathbf{34} \quad \int \frac{dx}{\cos x} = \int \sec x dx = \ln |\sec x + \tan x| + C; \int \frac{dx}{1+\cos x} \left(\frac{1-\cos x}{1-\cos x} \right) = \int \frac{dx}{\sin^2 x} - \int \frac{\cos x dx}{\sin^2 x} = \int \csc^2 x dx - \int \frac{du}{u^2} = -\cot x + \frac{1}{\sin x} = \frac{1-\cos x}{\sin x} + C; \int \frac{dx}{\sqrt{1+\cos x}} = \int \frac{dx}{\sqrt{2 \cos \frac{x}{2}}} = \sqrt{2} \ln \left| \sec \frac{x}{2} + \tan \frac{x}{2} \right| + C$$

$$\mathbf{36} \quad x = \tan \theta \text{ gives } \int \frac{dx}{\sqrt{x^2 + 1}} = \int \frac{\sec^2 \theta d\theta}{\sec \theta} = \ln (\sec \theta + \tan \theta) = \ln (x + \sqrt{x^2 + 1}) = g. \quad (\text{b}) \text{ Check } g' = \frac{1 + \frac{x}{\sqrt{x^2 + 1}}}{x + \sqrt{x^2 + 1}} = \frac{\sqrt{x^2 + 1} + x}{x + \sqrt{x^2 + 1}} \frac{1}{\sqrt{x^2 + 1}}. \text{ (c) Thus } \sinh g = \frac{1}{2} (e^g - e^{-g}) = \frac{1}{2} (x + \sqrt{x^2 + 1} - \frac{1}{x + \sqrt{x^2 + 1}}) = \frac{1}{2} \left(\frac{x^2 + 2x\sqrt{x^2 + 1} + x^2 + 1 - 1}{x + \sqrt{x^2 + 1}} \right) = x. \text{ (d) Now go directly to } \int \frac{dx}{\sqrt{x^2 + 1}} = \sinh^{-1} x \text{ by substituting } x = \sinh g \text{ to reach } \int \frac{\cosh g dg}{\cosh g} = g + C.$$

- 38 $x = \tanh \theta : \int \frac{dx}{x\sqrt{1-x^2}} = \int \frac{\operatorname{sech}^2 \theta d\theta}{\tanh \theta \operatorname{sech} \theta} = \int \operatorname{csch} \theta d\theta = -\ln |\operatorname{csch} \theta + \coth \theta| = -\ln(\frac{\sqrt{1-x^2+1}}{x}) + C$
- 40 $x = \cosh \theta : \int \frac{\sqrt{x^2-1}}{x^2} dx = \int \frac{\sinh \theta}{\cosh^2 \theta} \sinh \theta d\theta = \int \tanh^2 \theta d\theta = \int (1 - \operatorname{sech}^2 \theta) d\theta = \theta - \tanh \theta = \cosh^{-1} x - \frac{\sqrt{x^2-1}}{x} + C$
- 42 $x = \sinh \theta : \int \frac{\sqrt{1+x^2} dx}{x^2} = \int \frac{\cosh \theta}{\sinh^2 \theta} \cosh \theta d\theta = \int \coth^2 \theta d\theta = \int (1 + \operatorname{csch}^2 \theta) d\theta = \theta - \coth \theta = \sinh^{-1} x - \frac{\sqrt{x^2+1}}{x} + C$
- 44 $-x^2 + 2x + 8 = -(x-1)^2 + 9$ 46 $-x^2 + 10$: no linear term, square already completed
- 48 $x^2 + 4x - 12 = (x+2)^2 - 16$
- 50 $\int \frac{dx}{\sqrt{9-(x-1)^2}} = \int \frac{du}{\sqrt{9-u^2}}$. Set $u = 3 \sin \theta : \int \frac{\cos \theta d\theta}{\cos \theta} = \theta = \sin^{-1} \frac{u}{3} = \sin^{-1} \frac{x-1}{3} + C$;
 $\int \frac{dx}{10-x^2} = \frac{1}{2\sqrt{10}} \ln \frac{x-\sqrt{10}}{x+\sqrt{10}} + C$; $\int \frac{dx}{(x+2)^2-16} = \int \frac{du}{u^2-16} = \frac{1}{8} \ln \frac{2u-8}{2u+8} = \frac{1}{8} \ln \frac{x-2}{x+6} + C$
- 52 (a) $u = x-2$ (b) $u = x+1$ (c) $u = x-5$ (d) $u = x-\frac{1}{4}$
- 54 (a) If $x = \tan \theta$ then $\int \sqrt{1+x^2} dx = \int \sec^3 \theta d\theta$. (b) The integral $\frac{1}{2}[\sec \theta \tan \theta + \ln(\sec \theta + \tan \theta)]$ equals $\frac{1}{2}[x\sqrt{x^2+1} + \ln|x+\sqrt{x^2+1}|]$. (c) If $x = \sinh \theta$ then $\int \sqrt{1+x^2} dx = \int \cosh^2 \theta d\theta$ (d) The integral $\frac{1}{2}[\sinh \theta \cosh \theta + \theta]$ equals $\frac{1}{2}[x\sqrt{1+x^2} + \sinh^{-1} x]$.
- 56 The two curves cover the same area! Proof by calculus: $\int_0^4 \frac{dx}{\sqrt{16-x^2}} = (\text{with } x = 4u) \int_0^1 \frac{4du}{4\sqrt{1-u^2}}$. Proof by geometry: The x scale has factor $\frac{1}{4}$ and the y scale has factor 4, so $dA = dx dy$ is unchanged.

7.4 Partial Fractions (page 304)

The idea of partial fractions is to express $P(x)/Q(x)$ as a sum of simpler terms, each one easy to integrate. To begin, the degree of P should be less than the degree of Q . Then Q is split into linear factors like $x-5$ (possibly repeated) and quadratic factors like x^2+x+1 (possibly repeated). The quadratic factors have two complex roots, and do not allow real linear factors.

A factor like $x-5$ contributes a fraction $A/(x-5)$. Its integral is $A \ln(x-5)$. To compute A , cover up $x-5$ in the denominator of P/Q . Then set $x=5$, and the rest of P/Q becomes A . An equivalent method puts all fractions over a common denominator (which is Q). Then match the numerators. At the same point ($x=5$) this matching gives A .

A repeated linear factor $(x-5)^2$ contributes not only $A/(x-5)$ but also $B/(x-5)^2$. A quadratic factor like x^2+x+1 contributes a fraction $(Cx+D)/(x^2+x+1)$ involving C and D . A repeated quadratic factor or a triple linear factor would bring in $(Ex+F)/(x^2+x+1)^2$ or $G/(x-5)^3$. The conclusion is that any P/Q can be split into partial fractions, which can always be integrated.

- 1 $A = -1, B = 1, -\ln x + \ln(x-1) + C$ 3 $\frac{1}{x-3} - \frac{1}{x-2}$ 5 $\frac{1}{2x} - \frac{2}{x+1} + \frac{5/2}{x+2}$
 7 $\frac{3}{x} + \frac{1}{x^2}$ 9 $3 - \frac{3}{x^2+1}$ 11 $-\frac{1}{x} - \frac{1}{x^2} + \frac{1}{x-1}$ 13 $-\frac{1/6}{x} + \frac{1/2}{x-1} - \frac{1/2}{x-2} + \frac{1/6}{x-3}$
 15 $\frac{A}{x+1} + \frac{B}{x-1} + \frac{Cx+D}{x^2+1}; A = -\frac{1}{4}, B = \frac{1}{4}, C = 0, D = -\frac{1}{2}$
 17 Coefficients of $y: 0 = -Ab + B$; match constants $1 = Ac; A = \frac{1}{c}, B = \frac{b}{c}$
 19 $A = 1$, then $B = 2$ and $C = 1$; $\int \frac{dx}{x-1} + \int \frac{(2x+1)dx}{x^2+x+1} =$
 $\ln(x-1) + \ln(x^2+x+1) = \ln(x-1)(x^2+x+1) = \ln(x^3-1)$
 21 $u = e^x; \int \frac{du}{u^2-u} = \int \frac{du}{u-1} - \int \frac{du}{u} = \ln(\frac{u-1}{u}) + C = \ln(\frac{e^x-1}{e^x}) + C$

23 $u = \cos \theta; \int \frac{-du}{1-u^2} = -\frac{1}{2} \int \frac{du}{1-u} - \frac{1}{2} \int \frac{du}{1+u} = \frac{1}{2} \ln(1-u) - \frac{1}{2} \ln(1+u) = \frac{1}{2} \ln \frac{1-\cos \theta}{1+\cos \theta} + C$. We can reach $\frac{1}{2} \ln \frac{(1-\cos \theta)^2}{1-\cos^2 \theta} = \ln \frac{1-\cos \theta}{\sin \theta} = \ln(\csc \theta - \cot \theta)$ or a different way $\frac{1}{2} \ln \frac{1-\cos^2 \theta}{(1+\cos \theta)^2} = \ln \frac{\sin \theta}{1+\cos \theta} = -\ln \frac{1+\cos \theta}{\sin \theta} = -\ln(\csc \theta + \cot \theta)$

25 $u = e^x; du = e^x dx = u dx; \int \frac{1+u}{(1-u)u} du = \int \frac{2du}{1-u} + \int \frac{du}{u} = -2 \ln(1-e^x) + \ln e^x + C = -2 \ln(1-e^x) + x + C$

27 $x+1 = u^2, dx = 2u du; \int \frac{2u du}{1+u} = \int [2 - \frac{2}{1+u}] du = 2u - 2 \ln(1+u) + C = 2\sqrt{x+1} - 2 \ln(1+\sqrt{x+1}) + C$

29 Note $Q(a) = 0$. Then $\frac{x-a}{Q(x)} = \frac{x-a}{Q(x)-Q(a)} \rightarrow \frac{1}{Q'(a)}$ by definition of derivative. At a double root $Q'(a) = 0$.

2 $\frac{1}{(x-1)(x+1)} = \frac{A}{x-1} + \frac{B}{x+1}$. Cover up $x-1$ and set $x=1$ to find $A=\frac{1}{2}$. Cover up $x+1$ and set $x=-1$ to find $B=-\frac{1}{2}$. Then $\int \frac{dx}{x^2-1} = \frac{1}{2} \ln(x-1) - \frac{1}{2} \ln(x+1) = \frac{1}{2} \ln \frac{x-1}{x+1} + C$. Method 1: $\frac{1}{(x-1)(x+1)} = \frac{A(x+1)+B(x-1)}{(x-1)(x+1)}$ and by matching numerators $A+B=0$ and $A-B=1$ so again $A=\frac{1}{2}$ and $B=-\frac{1}{2}$.

$$\frac{x}{(x-3)(x-2)} = \frac{3}{x-3} - \frac{2}{x-2} \quad \frac{1}{x(x-1)(x+1)} = -\frac{1}{x} + \frac{1/2}{x-1} + \frac{1/2}{x+1}$$

$$\frac{3x+1}{(x-1)^2} = \frac{4}{(x-1)^2} + \frac{3}{x-1} \text{ (first multiply by } (x-1)^2 \text{ and set } x=1 \text{ to find the coefficient 4).}$$

$$\frac{1}{(x-1)(x^2+1)} = \frac{1/2}{x-1} - \frac{\frac{1}{2}x+\frac{1}{2}}{x^2+1} \quad \frac{x}{x^2-4} = \frac{1/2}{x-2} + \frac{1/2}{x+2}$$

$$x-1 + \frac{2}{x-1} \quad \text{so } \frac{x^2+1}{x+1} = x-1 + \frac{2}{x+1} \quad \frac{1}{x^2(x-1)} = -\frac{1}{x} - \frac{1}{x^2} + \frac{1}{x-1}$$

18 $\frac{x^2}{(x-3)(x+3)} = \frac{A(x+3)+B(x-3)}{(x-3)(x+3)}$ is impossible (no x^2 in the numerator on the right side).

Divide first to rewrite $\frac{x^2}{(x-3)(x+3)} = 1 + \frac{9}{(x-3)(x+3)} =$ (now use partial fractions) $1 + \frac{3/2}{x-3} - \frac{3/2}{x+3}$.

20 Integrate $\frac{1/2}{1-y} + \frac{1/2}{1+y}$ to find $-\frac{1}{2} \ln(1-y) + \frac{1}{2} \ln(1+y) = \frac{1}{2} \ln \frac{1+y}{1-y} = t + C$. At $t=0$ this is $\frac{1}{2} \ln 1 = 0 + C$ so $C=0$. Taking exponentials gives $\frac{1+y}{1-y} = e^{2t}$. Then $1+y = e^{2t}(1-y)$ and

$$y = \frac{e^{2t}-1}{e^{2t}+1} = \frac{e^t - e^{-t}}{e^t + e^{-t}} = \tanh t. \text{ This is the S-curve.}$$

22 Set $u = \sqrt{x}$ so $u^2 = x$ and $2u du = dx$. Then $\int \frac{1-\sqrt{x}}{1+\sqrt{x}} dx = \int \frac{1-u}{1+u} 2u du =$ (divide $u+1$ into $-2u^2 + 2u$) $\int (-2u+4 - \frac{4}{u+1}) du = -u^2 + 4u - 4 \ln(u+1) + C = -x + 4\sqrt{x} - 4 \ln(\sqrt{x}+1) + C$.

24 Set $u = e^t$ so $du = e^t dt$ or $dt = \frac{du}{u}$. Then $\int \frac{dt}{(e^t - e^{-t})^2} = \int \frac{du/u}{(u - \frac{1}{u})^2} = \int \frac{u du}{(u^2-1)^2} = \int (\frac{A}{u-1} + \frac{B}{(u-1)^2} + \frac{C}{u+1} + \frac{D}{(u+1)^2}) du$. Cover up $(u-1)^2$ and set $u=1$ to find $B=\frac{1}{4}$; cover up $(u+1)^2$ and set $u=-1$ to find $D=-\frac{1}{4}$; match left and right to find $A=C=0$. The integral is $-\frac{1}{4} \frac{1}{u-1} + \frac{1}{4} \frac{1}{u+1} = -\frac{1}{2} \frac{1}{u^2-1} = -\frac{1}{2} \frac{1}{e^{2t}-1}$. Check derivative: $\frac{1}{2} \frac{1}{(e^{2t}-1)^2} (2e^{2t}) = \frac{1}{(e^t - e^{-t})^2}$. Quicker integration: $\int \frac{u du}{(u^2-1)^2} = -\frac{1}{2} (u^2-1)^{-1} = -\frac{1}{2} \frac{1}{e^{2t}-1}$.

26 Set $u^3 = x-8$ so $3u^2 du = dx$. Then $\int \frac{(x-8)^{1/3} dx}{x} = \int \frac{u(3u^2 du)}{u^3+8} =$ (divide first) $\int (3 - \frac{24}{u^3+8}) du = 3u - \int \frac{24 du}{(u+2)(u^2-2u+4)} = 3u - \int (\frac{2}{u+2} + \frac{-2u+8}{u^2-2u+4}) du = 3u - 2 \ln(u+2) + \int \frac{2(u-1)-6}{(u-1)^2+3} du = 3u - 2 \ln(u+2) + \ln((u-1)^2+3) - \frac{6}{\sqrt{3}} \tan^{-1}(\frac{u-1}{\sqrt{3}}) + C$. Finally set $u = (x-8)^{1/3}$.

28 Set $u^4 = x$ so that $4u^3 du = dx$. Then $\int \frac{dx}{\sqrt{x} + \sqrt[4]{x}} = \int \frac{4u^3 du}{u^2+u} =$ (divide first) $\int (4u-4 + \frac{4u}{u^2+u}) du = 2u^2 - 4u + 4 \ln(u+1) + C = 2\sqrt{x} - 4\sqrt[4]{x} + 4 \ln(\sqrt[4]{x}+1) + C$.

30 Multiply $\frac{1}{x^8-1} = \frac{A}{x-1} + \dots$ by $x-1$ and let x approach 1 to find $A = \lim_{x \rightarrow 1} \frac{x-1}{x^8-1} = \lim_{x \rightarrow 1} \frac{1}{8x^7} = \frac{1}{8}$.

7.5 Improper Integrals (page 309)

An improper integral $\int_a^b y(x) dx$ has lower limit $a = -\infty$ or upper limit $b = \infty$ or y becomes infinite in the

interval $a \leq x \leq b$. The example $\int_1^\infty dx/x^3$ is improper because $b = \infty$. We should study the limit of $\int_1^b dx/x^3$ as $b \rightarrow \infty$. In practice we work directly with $-\frac{1}{2}x^{-2}|_1^\infty = \frac{1}{2}$. For $p > 1$ the improper integral $\int_0^1 x^{-p} dx$ is finite. For $p < 1$ the improper integral $\int_0^1 x^{-p} dx$ is finite. For $y = e^{-x}$ the integral from 0 to ∞ is 1.

Suppose $0 \leq u(x) \leq v(x)$ for all x . The convergence of $\int v(x) dx$ implies the convergence of $\int u(x) dx$. The divergence of $\int u(x) dx$ implies the divergence of $\int v(x) dx$. From $-\infty$ to ∞ , the integral of $1/(e^x + e^{-x})$ converges by comparison with $1/e^{|x|}$. Strictly speaking we split $(-\infty, \infty)$ into $(-\infty, 0)$ and $(0, \infty)$. Changing to $1/(e^x - e^{-x})$ gives divergence, because $e^x = e^{-x}$ at $x = 0$. Also $\int_{-\pi}^\pi dx/\sin x$ diverges by comparison with $\int dx/x$. The regions left and right of zero don't cancel because $\infty - \infty$ is not zero.

$$\begin{array}{lll} 1 \frac{x^{1-e}}{1-e}|_1^\infty = \frac{1}{e-1} & 3 -2(1-x)^{1/2}|_1^0 = 2 & 5 \tan^{-1} x|_{-\pi/2}^0 = \frac{\pi}{2} \\ 9 x \ln x - x|_0^e = -\infty & 11 \ln(\ln x)|_{100}^\infty = \infty & 13 \frac{1}{2}(x + \sin x \cos x)|_0^\infty = \infty \end{array}$$

- 15 $\frac{x^{1-p}}{1-p}|_0^\infty$ diverges for every p !
 17 Less than $\int_1^\infty \frac{dx}{x^6} = \frac{1}{5}$
 19 Less than $\int_0^1 \frac{dx}{x^2+1} + \int_1^\infty \frac{\sqrt{x} dx}{x^2} = \tan^{-1} x|_0^1 - \frac{2}{\sqrt{x}}|_1^\infty = \frac{\pi}{4} + 2$
 21 Less than $\int_1^\infty e^{-x} dx = \frac{1}{e}$, greater than $-\frac{1}{e}$
 23 Less than $\int_0^1 e^2 dx + e \int_1^\infty e^{-(x-1)^2} dx = e^2 + e \int_1^\infty e^{-u^2} du = e^2 + \frac{e}{\sqrt{\pi}}$
 25 $\int_0^1 \frac{\sin^2 x dx}{x^2} + \int_1^\infty \frac{\sin^2 x dx}{x^2}$ less than $1 + \int_1^\infty \frac{dx}{x^2} = 2$
 27 $p! = p$ times $(p-1)!$; 1 = 1 times 0!
 29 $u = x, dv = xe^{-x^2} dx : -x \frac{e^{-x^2}}{2}|_0^\infty + \int_0^\infty \frac{e^{-x^2}}{2} dx = \frac{1}{2}\sqrt{\pi}$
 31 $\int_0^\infty 1000e^{-1t} dt = -10,000e^{-1t}|_0^\infty = \$10,000$
 33 $W = \frac{-GMm}{x}|_R^\infty = \frac{GMm}{R} = \frac{1}{2}mv_0^2$ if $v_0 = \sqrt{\frac{2GM}{R}}$
 35 $\int_0^\infty \frac{dx}{2^x} = \int_0^\infty e^{-x \ln 2} dx = \frac{e^{-x \ln 2}}{-\ln 2}|_0^\infty = \frac{1}{\ln 2}$
 37 $\int_0^{\pi/2} (\sec x - \tan x) dx = [\ln(\sec x + \tan x) + \ln(\cos x)]_0^{\pi/2} = [\ln(1 + \sin x)]_0^{\pi/2} = \ln 2$.

The areas under $\sec x$ and $\tan x$ separately are infinite 39 Only $p = 0$

- 2 $\int_0^1 \frac{dx}{x^\pi} = [\frac{x^{1-\pi}}{1-\pi}]_0^1$ diverges at $x = 0$: infinite area 4 $\int_0^1 \frac{dx}{1-x} = [-\ln(1-x)]_0^1$ diverges at $x = 1$: infinite area
 6 $\int_{-1}^1 \frac{dx}{\sqrt{1-x^2}} = [\sin^{-1} x]_{-1}^1 = \frac{\pi}{2} - (-\frac{\pi}{2}) = \pi$
 8 $\int_{-\infty}^\infty \sin x dx$ is not defined because $\int_a^b \sin x dx = \cos a - \cos b$ does not approach a limit as $b \rightarrow \infty$
 and $a \rightarrow -\infty$
 10 $\int_0^\infty xe^{-x} dx = [-xe^{-x}]_0^\infty + \int_0^\infty e^{-x} dx = 0 + 1$
 12 $\int_{-\infty}^\infty \frac{x dx}{(x^2-1)^2}$ is not defined because the area around $x = -1$ and $x = 1$ is infinite.
 14 $\int_0^{\pi/2} \tan x dx$ is not defined: it is $\int_0^1 \frac{du}{u}$ with $u = \cos x$ and the area is infinite.
 16 $\int_0^\infty \frac{e^x dx}{(e^x-1)^p} = (\text{set } u = e^x - 1) \int_0^\infty \frac{du}{u^p}$ which is infinite: diverges at $u = 0$ if $p \geq 1$, diverges at $u = \infty$ if $p \leq 1$.
 18 $\int_0^1 \frac{dx}{x^2+1} < \int_0^1 \frac{dx}{1} = 1$: convergence 20 $\int_0^1 \frac{e^{-x} dx}{1-x} > \int_0^1 \frac{e^{-1} dx}{1-x} = \infty$: divergence
 22 $\int_1^\infty x^{-x} dx < \int_1^\infty e^{-x} dx = \frac{1}{e}$: convergence
 24 $\int_0^1 \sqrt{-\ln x} dx < \int_0^{1/e} (-\ln x) dx + \int_{1/e}^1 1 dx = [-x \ln x + x]|_0^{1/e} + [x]|_{1/e}^1 = \frac{1}{e} + 1$: convergence (note $x \ln x \rightarrow 0$ as $x \rightarrow 0$)
 26 $\int_0^\infty (\frac{1}{x} - \frac{1}{1+x}) dx$: the separate integrals would give $\infty - \infty$ which is indeterminate, so combine $\frac{1}{x} - \frac{1}{1+x} = \frac{1+x-x}{x(1+x)} < \frac{1}{x^2}$. The integral is less than $\int_1^\infty \frac{dx}{x^2} = 1$. Convergence.
 28 $\int_0^\infty x^{-1/2} e^{-x} dx$ (set $x = u^2$) = $\int_0^\infty u^{-1} e^{-u^2} 2u du = 2 \int_0^\infty e^{-u^2} du = \sqrt{\pi}$, so this is $(-\frac{1}{2})!$ Then $(p+1)! =$

$(p+1)$ times $p!$ with $p = -\frac{1}{2}$ gives $(\frac{1}{2})! = \frac{1}{2}\sqrt{\pi}$.

30 $B(m, n) = \int_0^1 x^{m-1}(1-x)^{n-1} dx$ is like $\int x^{m-1} dx$ near $x=0$ and $\int(1-x)^{n-1} dx$ near $x=1$. These are finite if $m-1 > -1$ and $n-1 > -1$, or $m > 0$ and $n > 0$. Then the front inside cover gives $B = \frac{(m-1)!(n-1)!}{(m+n-1)!}$.

32 To pay s at the end of year n , the present deposit must be $\frac{s}{(1+i)^n} = \frac{s}{a^n}$. To pay s at the end of every year (perpetual annuity), the deposit must be $\frac{s}{a} + \frac{s}{a^2} + \dots = s \frac{1/a}{1-1/a} = \frac{s}{a-1} = \frac{s}{i}$. To receive $s = \$1000/\text{year}$ with $i = 10\%$ you deposit \$10,000.

34 Note: $GM = 4 \cdot 10^{14} \text{ m}^3/\text{sec}^2$: the lost factor of 10^{10} would have a large effect on our universe! The escape velocity is $v_0 = \sqrt{2GM/R}$, so that $R = 2GM/v_0^2 = 2 \cdot 4 \cdot 10^{14}/9 \cdot 10^{16} = \frac{8}{9} \cdot 10^{-2}$ meters = .9 cm.

36 $\int_a^b \frac{x \, dx}{1+x^2} = [\frac{1}{2} \ln(1+x^2)]_a^b = \frac{1}{2} \ln(1+b^2) - \frac{1}{2} \ln(1+a^2)$. As $b \rightarrow \infty$ or as $a \rightarrow -\infty$ (separately!) there is no limiting value. If $a = -b$ then the answer is zero – but we are not allowed to connect a and b .

38 $\int_0^\infty \frac{x^{-1/2} \, dx}{1+x} = (\text{set } x = u^2) \int_0^\infty \frac{(\frac{1}{2})2u \, du}{1+u^2} = [2 \tan^{-1} u]_0^\infty = 2(\frac{\pi}{2}) = \pi$; $\int_0^\infty xe^{-x} \cos x \, dx = (\text{by parts})$
 $[\frac{xe^{-x}}{2}(\sin x - \cos x) + \frac{e^{-x}}{2} \sin x]_0^\infty = 0$.

40 The red area in the right figure has an extra unit square (area 1) compared to the red area on the left.

MIT OpenCourseWare
<https://ocw.mit.edu>

Resource: Calculus
Gilbert Strang

The following may not correspond to a particular course on MIT OpenCourseWare, but has been provided by the author as an individual learning resource.

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.